


# Imprecision in Martingale-Theoretic Prequential Randomness

by Floris Persiau and Gert de Cooman






# Imprecision in Martingale-Theoretic Prequential Randomness

by Floris Persiau and Gert de Cooman


# Imprecision in Martingale-Theoretic Prequential Randomness



$$\leftrightarrow 1/2$$



# Imprecision in Martingale-Theoretic Prequential Randomness




$$\leftrightarrow 1/2$$

0101010101...



# Imprecision in Martingale-Theoretic Prequential Randomness



$$\leftrightarrow 1/2$$

1010010011...




# Imprecision in Martingale-Theoretic Prequential Randomness

Path(s)

$$\omega \in \{0, 1\}^{\mathbb{N}}$$

Situation(s)

$$w \in \{0, 1\}^* := \bigcup_{n \in \mathbb{N}_0} \{0, 1\}^n$$



$$\leftrightarrow 1/2$$

1010010011...






# Imprecision in Martingale-Theoretic Prequential Randomness

Path(s)

$$\omega \in \{0, 1\}^{\mathbb{N}}$$

Situation(s)

$$w \in \{0, 1\}^* := \bigcup_{n \in \mathbb{N}_0} \{0, 1\}^n$$



$$\leftrightarrow 1/2$$

1010010011...



# Imprecision in Martingale-Theoretic Prequential Randomness

Path(s)


$$\omega \in \{0, 1\}^{\mathbb{N}}$$

Situation(s)

$$w \in \{0, 1\}^* := \bigcup_{n \in \mathbb{N}_0} \{0, 1\}^n$$

Forecasting system

$$\varphi: \{0, 1\}^* \rightarrow [0, 1]$$




$$\leftrightarrow 1/2$$



1010010011...

# Imprecision in Martingale-Theoretic Prequential Randomness



# Imprecision in Martingale-Theoretic Prequential Randomness



Martin-Löf  
randomness



Computable  
randomness



Church  
randomness



weak Martin-Löf  
randomness




Schnorr  
randomness




weak Church  
randomness

# Imprecision in Martingale-Theoretic Prequential Randomness



Martin-Löf  
randomness



Computable  
randomness



Church  
randomness



weak Martin-Löf  
randomness



Schnorr  
randomness



weak Church  
randomness

# Imprecision in Martingale-Theoretic **Prequential** Randomness

$$\omega \leftrightarrow \varphi$$


## Betting strategies

A real-valued map  $T: \{0,1\}^* \rightarrow \mathbb{R}$  is called a **test supermartingale** for a forecasting system  $\varphi$  if



- i)  $T(\square) = 1$ ,
- ii)  $T(w) \geq 0$  for all  $w \in \{0,1\}^*$ ,
- iii)  $\bar{E}_{\varphi(w)}(T(w \cdot)) \leq T(w)$  for all  $w \in \{0,1\}^*$ .

## Definition

A path  $\omega \in \{0,1\}^{\mathbb{N}}$  is **Martin-Löf random** for a forecasting system  $\varphi$  if  $\limsup_{n \rightarrow +\infty} T(\omega_1, \dots, \omega_n) < +\infty$  for all lower semicomputable test supermartingales  $T$  for  $\varphi$ .





# Imprecision in Martingale-Theoretic **Prequential** Randomness



€  $\leftrightarrow$  1/2



# Imprecision in Martingale-Theoretic Prequential Randomness




€  $\leftrightarrow$  1/2






# Imprecision in Martingale-Theoretic Prequential Randomness



€ ↔ 1/2



Preq. path(s)

$$(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_r \times \{0, 1\})^{\mathbb{N}}$$


Preq. situation(s)

$$(I_1, \omega_1, \dots, I_n, \omega_n) \in (\mathcal{I}_r \times \{0, 1\})^*$$

? ? ?



# Imprecision in Martingale-Theoretic Prequential Randomness




## Betting strategies

A real-valued map  $T: \{0,1\}^* \rightarrow \mathbb{R}$  is called a **test supermartingale** for a forecasting system  $\varphi$  if

- i)  $T(\square) = 1$ ,
- ii)  $T(w) \geq 0$  for all  $w \in \{0,1\}^*$ ,
- iii)  $\bar{E}_{\varphi(w)}(T(w \cdot)) \leq T(w)$  for all  $w \in \{0,1\}^*$ .

## Definition

A path  $\omega \in \{0,1\}^{\mathbb{N}}$  is **Martin-Löf random** for a forecasting system  $\varphi$  if  $\limsup_{n \rightarrow +\infty} T(\omega_1, \dots, \omega_n) < +\infty$  for all lower semicomputable test supermartingales  $T$  for  $\varphi$ .



## Betting strategies


A real-valued map  $F: (\mathcal{I}_{\mathbb{R}} \times \{0,1\})^* \rightarrow \mathbb{R}$  is called a **test superfarthingale** if

- i)  $F(\square) = 1$ ,
- ii)  $F(I_1, \omega_1, \dots, I_n, \omega_n) \geq 0$  for all  $(I_1, \omega_1, \dots, I_n, \omega_n) \in (\mathcal{I}_{\mathbb{R}} \times \{0,1\})^*$ ,
- iii)  $\bar{E}_{I_r}(F(I_1, \omega_1, \dots, I_n, \omega_n, I_r, \cdot)) \leq F(I_1, \omega_1, \dots, I_n, \omega_n)$  for all  $(I_1, \omega_1, \dots, I_n, \omega_n) \in (\mathcal{I}_{\mathbb{R}} \times \{0,1\})^*$  and  $I_r \in \mathcal{I}_{\mathbb{R}}$ .

## Definition

We call a sequence  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_{\mathbb{R}} \times \{0,1\})^{\mathbb{N}}$  of rational forecasts and outcomes **game-random** if it's non-degenerate and if  $\limsup_{n \rightarrow +\infty} F(I_1, \omega_1, \dots, I_n, \omega_n) < +\infty$  for all lower semicomputable test superfarthingales  $F$ .

# Imprecision in Martingale-Theoretic Prequential Randomness




## Betting strategies

A real-valued map  $T: \{0,1\}^* \rightarrow \mathbb{R}$  is called a **test supermartingale** for a forecasting system  $\varphi$  if

- i)  $T(\square) = 1$ ,
- ii)  $T(w) \geq 0$  for all  $w \in \{0,1\}^*$ ,
- iii)  $\bar{E}_{\varphi(w)}(T(w \cdot)) \leq T(w)$  for all  $w \in \{0,1\}^*$ .

## Definition

A path  $\omega \in \{0,1\}^{\mathbb{N}}$  is **Martin-Löf random** for a forecasting system  $\varphi$  if  $\limsup_{n \rightarrow +\infty} T(\omega_1, \dots, \omega_n) < +\infty$  for all lower semicomputable test supermartingales  $T$  for  $\varphi$ .



## Betting strategies


A real-valued map  $F: (\mathcal{I}_{\mathbf{r}} \times \{0,1\})^* \rightarrow \mathbb{R}$  is called a **test superfarthingale** if

- i)  $F(\square) = 1$ ,
- ii)  $F(I_1, \omega_1, \dots, I_n, \omega_n) \geq 0$  for all  $(I_1, \omega_1, \dots, I_n, \omega_n) \in (\mathcal{I}_{\mathbf{r}} \times \{0,1\})^*$ ,
- iii)  $\bar{E}_{I_{\mathbf{r}}}(F(I_1, \omega_1, \dots, I_n, \omega_n, I_{\mathbf{r}}, \cdot)) \leq F(I_1, \omega_1, \dots, I_n, \omega_n)$  for all  $(I_1, \omega_1, \dots, I_n, \omega_n) \in (\mathcal{I}_{\mathbf{r}} \times \{0,1\})^*$  and  $I_{\mathbf{r}} \in \mathcal{I}_{\mathbf{r}}$ .

## Definition

We call a sequence  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_{\mathbf{r}} \times \{0,1\})^{\mathbb{N}}$  of **rational** forecasts and outcomes **game-random** if it's non-degenerate and if  $\limsup_{n \rightarrow +\infty} F(I_1, \omega_1, \dots, I_n, \omega_n) < +\infty$  for all lower semicomputable test superfarthingales  $F$ .

# Imprecision in Martingale-Theoretic Prequential Randomness




## Rational restriction

For every non-degenerate computable forecasting system  $\varphi$  there's a recursive rational forecasting system  $\varphi_r$  such that a path  $\omega \in \{0, 1\}^{\mathbb{N}}$  is Martin-Löf random for  $\varphi$  if and only if it's Martin-Löf random for  $\varphi_r$ .

### Definition

A path  $\omega \in \{0, 1\}^{\mathbb{N}}$  is **Martin-Löf random** for a forecasting system  $\varphi$  if  $\limsup_{n \rightarrow +\infty} T(\omega_1, \dots, \omega_n) < +\infty$  for all lower semicomputable test supermartingales  $T$  for  $\varphi$ .



## Betting strategies


A real-valued map  $F: (\mathcal{I}_r \times \{0, 1\})^* \rightarrow \mathbb{R}$  is called a **test superfarthingale** if

- i)  $F(\square) = 1$ ,
- ii)  $F(I_1, \omega_1, \dots, I_n, \omega_n) \geq 0$  for all  $(I_1, \omega_1, \dots, I_n, \omega_n) \in (\mathcal{I}_r \times \{0, 1\})^*$ ,
- iii)  $\bar{E}_{I_r}(F(I_1, \omega_1, \dots, I_n, \omega_n, I_r, \cdot)) \leq F(I_1, \omega_1, \dots, I_n, \omega_n)$  for all  $(I_1, \omega_1, \dots, I_n, \omega_n) \in (\mathcal{I}_r \times \{0, 1\})^*$  and  $I_r \in \mathcal{I}_r$ .

## Definition

We call a sequence  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_r \times \{0, 1\})^{\mathbb{N}}$  of **rational** forecasts and outcomes **game-random** if it's non-degenerate and if  $\limsup_{n \rightarrow +\infty} F(I_1, \omega_1, \dots, I_n, \omega_n) < +\infty$  for all lower semicomputable test superfarthingales  $F$ .

# Imprecision in Martingale-Theoretic Prequential Randomness



compatible if  $(\forall n \in \mathbb{N}_0) \varphi_r(\omega_1, \dots, \omega_n) = I_{n+1}$

## Martin-Löf '=' Prequential

Consider any non-degenerate recursive rational forecasting system  $\varphi_r$ . Then any path  $\omega \in \{0,1\}^{\mathbb{N}}$  is Martin-Löf random for  $\varphi_r$  if and only if the prequential path  $(\varphi_r(\square), \omega_1, \varphi_r(\omega_1), \omega_2, \dots) \in (\mathcal{I}_r \times \{0,1\})^{\mathbb{N}}$  is game-random.




# Imprecision in Martingale-Theoretic Sequential Randomness

Consider a fair coin—so with probability 1/2 for heads—and assume that we toss it an infinite number of times. For this classical set-up, you are familiar with statements as 'The set of sequences that satisfy the law of large numbers has measure one w.r.t. the obvious induced measure'. You are probably less acquainted with examining the properties of a single sequence that is generated by tossing this coin. In Algorithmic Randomness, we study what it means for such infinite sequences to be 'random' w.r.t. 1/2. When adopting a martingale-theoretic approach, a sequence is then basically considered to be random if you cannot get arbitrarily rich by betting on the successive outcomes.

Classical martingale-theoretic notions of randomness not only allow for defining an infinite sequence's randomness w.r.t. probability 1/2, but also w.r.t. precise forecasting systems, which are maps from binary strings to probabilities. However, it is only rather recently that we succeeded at allowing for interval-valued forecasting systems in such notions of randomness. By doing so, we can test whether an infinite binary sequence is random w.r.t. a set of so-called compatible precise forecasting systems.

It is not always natural to define an infinite sequence's randomness w.r.t. an (imprecise) forecasting system. Consider for example a weather forecaster who provides a daily probability interval for rain in the next 24 hours. He typically only specifies forecasts 'on the fly', that is, based on the rain history he has actually observed, and doesn't provide forecasts for all rain histories that might have been and might be. As a way out of this conundrum, we adopt Dawid and Vovk's *prequential forecasting* framework, which allows us to define what it means for an infinite sequence  $(I_1, x_1, I_2, x_2, \dots)$  of successive rational interval forecasts  $I_i$  and subsequent binary outcomes  $x_i$  to be random.

interval forecast(s)  $I \in \mathcal{I} := \{[p, q] \subseteq [0, 1] : p \leq q \in \mathbb{R}\}$   
 situation(s)  $w \in \{0, 1\}^* := \bigcup_{n \in \mathbb{N}_0} \{0, 1\}^n$   
 path(s)  $\omega \in \{0, 1\}^{\mathbb{N}}$   
 (rat.) forecasting system  $\varphi_{(I)}: \{0, 1\}^* \rightarrow \mathcal{I}_{(I)}$   
*non-degenerate* if  $(\forall w \in \{0, 1\}^*) \varphi_{(I)}(w) \not\subseteq \{0, 1\}$



rat. interval forecast(s)  $I_i \in \mathcal{I}_i := \{[p, q] \subseteq [0, 1] : p \leq q \in \mathbb{Q}\}$   
 preq. situation(s)  $(I_1, \omega_1, \dots, I_n, \omega_n) \in (\mathcal{I}_i \times \{0, 1\})^n$   
 preq. path(s)  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_i \times \{0, 1\})^{\mathbb{N}}$   
*non-degenerate* if  $(\forall n \in \mathbb{N}) I_n \not\subseteq \{1 - \omega_n\}$

## Martin-Löf randomness

**Betting strategies**  
 A real-valued map  $T: \{0, 1\}^* \rightarrow \mathbb{R}$  is called a **test supermartingale** for a forecasting system  $\varphi$  if  $T(\square) = 1$ ,  $T(w) \geq 0$  and  $E_{\varphi(w)}(T(w \cdot)) \leq T(w)$  for all  $w \in \{0, 1\}^*$ .

**Definition**  
 A path  $\omega \in \{0, 1\}^{\mathbb{N}}$  is **Martin-Löf random** for a forecasting system  $\varphi$  if  $\limsup_{n \rightarrow \infty} T(\omega_1, \dots, \omega_n) < +\infty$  for all lower semicomputable test supermartingales  $T$  for  $\varphi$ .

## Rational restriction

For every non-degenerate computable forecasting system  $\varphi$  there's a recursive rational forecasting system  $\varphi_r$  with  $\varphi_r \subseteq \varphi$ , such that a path  $\omega \in \{0, 1\}^{\mathbb{N}}$  is Martin-Löf random for  $\varphi$  if and only if it's Martin-Löf random for  $\varphi_r$ .

**Property 1: universal supermartingale**  
 For every non-degenerate computable forecasting system  $\varphi$ , there's a so-called *universal* lower semicomputable test supermartingale  $T$  such that any path  $\omega \in \{0, 1\}^{\mathbb{N}}$  is Martin-Löf random for  $\varphi$  if and only if  $\lim_{n \rightarrow \infty} T(\omega_1, \dots, \omega_n) < +\infty$ .

**Property 2: at least one random path**  
 For every forecasting system  $\varphi$ , there is at least one path  $\omega \in \{0, 1\}^{\mathbb{N}}$  that is Martin-Löf random for  $\varphi$ .

**Property 3: monotonicity**  
 Consider any forecasting system  $\varphi$ . If a path  $\omega \in \{0, 1\}^{\mathbb{N}}$  is Martin-Löf random for  $\varphi$ , then it is Martin-Löf random as well for any forecasting system  $\psi$  for which  $\varphi \subseteq \psi$ .

**Property 4: relative frequencies**  
 Consider any path  $\omega \in \{0, 1\}^{\mathbb{N}}$ , any computable forecasting system  $\varphi$  and any recursive selection process  $S: \{0, 1\}^* \rightarrow \{0, 1\}$  such that  $\lim_{n \rightarrow \infty} \sum_{i=1}^n S(\omega_1, \dots, \omega_i) = +\infty$ . If  $\omega$  is Martin-Löf random for  $\varphi$ , then


$$\liminf_{n \rightarrow \infty} \frac{\sum_{i=1}^n S(\omega_1, \dots, \omega_i) [\omega_{i+1} - \min \varphi(\omega_1, \dots, \omega_i)]}{\sum_{i=1}^n S(\omega_1, \dots, \omega_i)} \geq 0$$

and


$$\limsup_{n \rightarrow \infty} \frac{\sum_{i=1}^n S(\omega_1, \dots, \omega_i) [\omega_{i+1} - \max \varphi(\omega_1, \dots, \omega_i)]}{\sum_{i=1}^n S(\omega_1, \dots, \omega_i)} \leq 0.$$

## Martin-Löf '=<sup>' Prequential</sup>

Consider any non-degenerate recursive rational forecasting system  $\varphi_r$ . Then any path  $\omega \in \{0, 1\}^{\mathbb{N}}$  is Martin-Löf random for  $\varphi_r$  if and only if the prequential path  $(\varphi_r(\square), \omega, \varphi_r(\omega_1), \omega_2, \dots) \in (\mathcal{I}_i \times \{0, 1\})^{\mathbb{N}}$  is game-random.



QR code: What kind of forecasting framework do we borrow from Dawid and Vovk?



Floris Persiau & Gert de Cooman

## Prequential randomness

**Betting strategies**  
 A real-valued map  $F: (\mathcal{I}_i \times \{0, 1\})^* \rightarrow \mathbb{R}$  is called a **test superfairthingale** if  $F(\square) = 1$ ,  $F(I_1, \omega_1, \dots, I_n, \omega_n) \geq 0$  and  $E_{I_i}(F(I_1, \omega_1, \dots, I_n, \omega_n, I_{n+1})) \leq F(I_1, \omega_1, \dots, I_n, \omega_n)$  for all  $(I_1, \omega_1, \dots, I_n, \omega_n) \in (\mathcal{I}_i \times \{0, 1\})^n$  and  $I_{n+1} \in \mathcal{I}_i$ .

**Definition**  
 We call a sequence  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_i \times \{0, 1\})^{\mathbb{N}}$  of rational forecasts and outcomes **game-random** if it's non-degenerate and if all lower semicomputable test superfairthingales  $F$  satisfy  $\limsup_{n \rightarrow \infty} F(I_1, \omega_1, \dots, I_n, \omega_n) < +\infty$ .

## Martin-Löf $\subseteq$ Prequential

Consider any infinite sequence of interval forecasts and outcomes  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_i \times \{0, 1\})^{\mathbb{N}}$  that's game-random. Then the infinite sequence of outcomes  $\omega$  is Martin-Löf random for any rational forecasting system  $\varphi_r$  that's compatible with  $(I_1, \omega_1, I_2, \omega_2, \dots)$ .

**Property 1: universal superfairthingale**  
 There's a so-called *universal* lower semicomputable test superfairthingale  $U$  with the property that any non-degenerate prequential path  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_i \times \{0, 1\})^{\mathbb{N}}$  is game-random if and only if  $\limsup_{n \rightarrow \infty} U(I_1, \omega_1, \dots, I_n, \omega_n) < +\infty$ .

**Property 2: at least one random preq. path**  
 For every infinite sequence of rational interval forecasts  $(I_1, I_2, \dots) \in \mathcal{I}_i^{\mathbb{N}}$  there's at least one path  $\omega \in \{0, 1\}^{\mathbb{N}}$  such that  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_i \times \{0, 1\})^{\mathbb{N}}$  is game-random.

**Property 3: monotonicity**  
 Consider any recursive rational forecasting system  $\varphi_r$  and any game-random prequential path  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_i \times \{0, 1\})^{\mathbb{N}}$ . If  $I_{n+1} \subseteq \varphi_r(\omega_1, \dots, \omega_n)$  for all  $n \in \mathbb{N}_0$ , then  $(\varphi_r(\square), \omega, \varphi_r(\omega_1), \omega_2, \dots) \in (\mathcal{I}_i \times \{0, 1\})^{\mathbb{N}}$  is game-random as well.

**Property 4: relative frequencies**  
 Consider any infinite sequence of rational interval forecasts and outcomes  $(I_1, \omega_1, I_2, \omega_2, \dots) \in (\mathcal{I}_i \times \{0, 1\})^{\mathbb{N}}$  and any recursive selection function  $S: (\mathcal{I}_i \times \{0, 1\})^* \times \mathcal{I}_i \rightarrow \{0, 1\}$  such that  $\sum_{i=1}^n S(I_1, \omega_1, \dots, I_i, \omega_i, I_{i+1}) = +\infty$ . If  $(I_1, \omega_1, I_2, \omega_2, \dots)$  is game-random, then

$$\liminf_{n \rightarrow \infty} \frac{\sum_{i=1}^n S(I_1, \omega_1, \dots, I_i, \omega_i, I_{i+1}) [\omega_{i+1} - \min I_{i+1}]}{\sum_{i=1}^n S(I_1, \omega_1, \dots, I_i, \omega_i, I_{i+1})} \geq 0$$

and

$$\limsup_{n \rightarrow \infty} \frac{\sum_{i=1}^n S(I_1, \omega_1, \dots, I_i, \omega_i, I_{i+1}) [\omega_{i+1} - \max I_{i+1}]}{\sum_{i=1}^n S(I_1, \omega_1, \dots, I_i, \omega_i, I_{i+1})} \leq 0.$$
